QUANTIZATION OF THE ELECTRIC CHARGE OF LEPTONS

B.I. MEHDIYEV
Institute of Physics, Ministry of Science and Education of Azerbaijan, AZ-1073, Baku, Azerbaijan
bahram.mehdiyev@mail.ru

The possibility of obtaining electric charge quantization for leptons within the Weinberg-Salam model is investigated for the case where the right-handed electron component is included in an isodoublet. Interaction Lagrangians describing the coupling of leptons and Higgs fields to gauge fields were derived. In doing so, specific values for the weak hypercharges were not assumed; instead, they were treated as free parameters to be determined based on physical requirements. Relations derived from anomaly cancellation conditions have not been utilized. It is shown that the interaction of the Higgs field with lepton isomultiplets leads to specific relations between the hypercharges of these fields. Taking into account the conservation of hypercharge and the P-invariance of the electromagnetic interaction between the electron and the new lepton (E^0), an expression for the electric charge of the leptons has been obtained that depends solely on the hypercharge of the left-handed lepton isodoublet and the Higgs isodoublet. This can be viewed as evidence of the quantization of the electric charge of leptons. The P-invariance of the electromagnetic interactions of the introduced leptons implies that the photon mass is zero (MA=0)), that the masses of the $W^{\pm}$ and Z bosons satisfy the known relation $M_W = M_Z \cos\theta$, and that the resulting values for the lepton charges are consistent with experimental data.

Keywords: lepton isodoublet, Higgs fields, hypercharge, P-invariance, quantization condition.
DOI: https://doi.org/10.70784/ajp.1.2026103

INTRODUCTION

A number of papers [1-11] was dedicated to the study of the quantization of electric charge in the Standard Model (SM) and its extensions. In [12-14], we investigated the quantization of the electric charge of leptons and quarks in the SM. Unlike [1-11], in [12-14] did not use relations that follow from the anomaly cancellation conditions. In [12, 14], the right-hand component of the electron is introduced as an isosinglet e_R^-. In this paper, we introduce the right-hand component of the electron as a component of isodoublet $\begin{pmatrix} E^0 \\ e^- \end{pmatrix}_R$ (E^0 - a new lepton). As in [12-14], here either we will not use relations that follow from the anomaly cancellation conditions.

Thus, in this case we have the following lepton fields:

$$\psi_{eL} = \begin{pmatrix} \nu_e(\alpha) \\ e^- \end{pmatrix}_L ,$$

$$\psi_{eR} = \begin{pmatrix} E^0 \\ e^- \end{pmatrix}_R ,$$

$$\psi_L^{'} = E_L^0(\alpha) \tag{1}$$

and the Higgs isodoublet φ and isosinglet ϕ:

$$\varphi = \begin{pmatrix} \varphi_+ \\ \varphi_0 \end{pmatrix}, \phi ; \tag{2}$$

where

$$\nu_{eL}(\alpha) = \nu_{eL} cos\alpha + E_L^0 sin\alpha,$$
$$E_L^0(\alpha) = E_L^0 cos\alpha - \nu_{eL} sin\alpha,$$

(we introduce mixing ν_{eL} and E_L^0 to eliminate off-diagonal terms from the mass part of the Lagrangian). The Lagrangian for the interaction of leptons and Higgs fields with gauge fields is obtained by replacing

$$\partial_\mu \to D_\mu = \partial_\mu - ig\left(\vec{t}\vec{b}_\mu\right) - i\frac{Y}{2} g_1 a_\mu \tag{3}$$

in the Lagrangian of the Higgs field and the free massless Dirac field $i\overline{\psi}\gamma_\mu\partial_\mu\psi$, and the desired Lagrangian has the form

$$L = i\overline{\psi}_{eL}\gamma_\mu D_\mu^L \psi_{eL} + i\overline{\psi}_{eR}\gamma_\mu D_\mu^R \psi_{eR} + i\psi_L^{'}\gamma_\mu D_\mu^{L'}\psi_L^{'} + \left(D_\mu^\varphi\varphi\right)^+\left(D_\mu^\varphi\varphi\right) + \left(D_\mu^\phi\phi\right)^+\left(D_\mu^\phi\phi\right) \tag{4}$$

Here $g, g_1 -$ are the interaction constants, $\vec{t}$ and $Y -$ is the isospin and hypercharge operator of a given multiplet, $\vec{b}_\mu -$ is the Yang-Mills isotriplet, $a_\mu -$ is the Maxwell isosinglet; ψ_{eL}, ψ_{eR} and $\psi_L^{'} -$ are the left and right isomultiplets of the lepton fields. We do not accept the Gell-Mann–Nishijima relation for weak hypercharges. We will accept them as free parameters and will try to fix them based on physical requirements.

We denote the hypercharges of the lepton and Higgs fields as follows

$$Y\left(\psi_{eL}\right) = y_L, \quad Y\left(\psi_{eR}\right) = y_R, \quad Y\left(\psi_L^{'}\right) = y_1, \qquad Y(\varphi) = y_\varphi, Y(\phi) = y_\phi \tag{5}$$

131, H.Javid ave, AZ-1073, Baku
Institute of Physics
E-mail: jophphysics@gmail.com

and assume they are real.

As is well known, for isoscalar fields $E_L^{'}$ and ϕ: $\vec{t}=0$, and for isodoublets ψ_{eL},ψ_{eR} and φ: $\vec{t}=\vec{\tau}/2$ ($\vec{\tau}-$ Pauli matrices). We write the transformation of the fields b_μ^3 and a_μ into physical fields A_μ and Z_μ as

$$\begin{aligned} b_\mu^3 &= A_\mu \sin\theta + Z_\mu \cos\theta, \\ a_\mu &= A_\mu \cos\theta - Z_\mu \sin\theta, \end{aligned} \qquad (6)$$

where $\sin\theta=\frac{g_1}{\sqrt{g^2+g_1^2}}$, $\cos\theta=\frac{g}{\sqrt{g^2+g_1^2}}$.

THE INTERACTION OF LEPTONS WITH VECTOR BOSONS

Taking into account (5) in (4), for the interaction of leptons with gauge fields, we have

$$\begin{aligned} L_l &= \overline{\psi}_{eL}(i\gamma_\mu D_\mu^L)\psi_{eL} + \overline{\psi}_{eR}(i\gamma_\mu D_\mu^R)\psi_{eR} + \overline{\psi}_L^{'}(i\gamma_\mu D_\mu^{L^{'}})\psi_L^{'} = \\ &= \overline{\psi}_{eL}\gamma_\mu[i\partial_\mu + \frac{1}{2}(g\vec{\tau}\vec{b}_\mu + g_1 y_L a_\mu)]\psi_{eL} + \overline{\psi}_{eR}\gamma_\mu\left[i\partial_\mu + \frac{1}{2}(g\vec{\tau}\vec{b}_\mu + g_1 y_R a_\mu)\right]\psi_{eR} + \\ &+ \overline{\psi}_L^{'}\gamma_\mu(i\partial_\mu + \frac{1}{2}g_1 y_1 a_\mu)\psi_L^{'} \equiv L_{KIN} + L_{CC} + L_{NC}. \end{aligned} \qquad (7)$$

Here $L_{KIN} = i\overline{\nu}_{eL}(\alpha)\hat{\partial}\nu_{eL}(\alpha) + i\overline{e}_L^-\hat{\partial}e_L^- + i\overline{e}_R^-\hat{\partial}e_R^- + i\overline{E}_R^0\hat{\partial}E_R^0 + i\overline{E}_L^0(\alpha)\hat{\partial}E_L^0(\alpha) -$
is the kinetic part of the Lagrangian,

$$L_{CC} = \frac{g}{\sqrt{2}}\left[(\overline{\nu}_{eL}(\alpha)\gamma_\mu e_L^- + \overline{E}_R^0\gamma_\mu e_R^-)W_\mu^+ + (\overline{e}_L^-\gamma_\mu\nu_{eL}(\alpha) + \overline{e}_R^-\gamma_\mu E_R^0)W_\mu^-\right] -$$

is the Lagrangian of the interaction of charged currents (CC),

$$\begin{aligned} L_{NC} &= \frac{1}{2}\overline{\nu}_{eL}(\alpha)\gamma_\mu\nu_{eL}(\alpha)(gb_\mu^3 + g_1 y_L a_\mu) + \frac{1}{2}\overline{e}_L\gamma_\mu e_L(-gb_\mu^3 + g_1 y_L a_\mu) + \frac{1}{2}\overline{e}_R\gamma_\mu e_R(-gb_\mu^3 + g_1 y_R a_\mu) + \\ &+ \frac{1}{2}\overline{E}_R^0\gamma_\mu E_R^0(gb_\mu^3 + g_1 y_R a_\mu) + \frac{1}{2}\overline{E}_L^0(\alpha)\gamma_\mu E_L^0(\alpha)g_1 y_1 a_\mu - \end{aligned}$$

is the Lagrangian of the interaction of neutral currents (NC).

Taking into account the relations $\bar{f}_L\gamma_\mu\psi_L = \frac{1}{2}\bar{f}O_\mu\psi$, $\bar{f}_R\gamma_\mu\psi_R = \frac{1}{2}\bar{f}O_\mu^{'}\psi$ and moving on to the physical fields A_μ and Z_μ (relationship (6)), for the parts of the Lagrangians L_{CC} and L_{NC}, we have the following expressions:

$$L_{CC} = \frac{g}{2\sqrt{2}}\left[(\overline{\nu}_{eL}(\alpha)O_\mu e^- + \overline{E}^0 O_\mu^{'} e^-)W_\mu^+ + (\overline{e}^- O_\mu\nu_e(\alpha) + \overline{e}^- O_\mu^{'}E^0)W_\mu^-\right] \qquad (8)$$

$$\begin{aligned} L_{NC} &= \frac{1}{4}\overline{\nu}_e(\alpha)O_\mu\nu_e(\alpha)(gb_\mu^3 + g_1 y_L a_\mu) + \frac{1}{4}\overline{e}^-\gamma_\mu\left[-2gb_\mu^3 + g_1(y_L + y_R)a_\mu + \gamma_5 g_1(y_L - y_R)a_\mu\right]e^- + \\ &+ \frac{1}{4}\overline{E}^0\gamma_\mu\left[gb_\mu^3 + g_1(y_1 + y_R)a_\mu + \gamma_5\left(-gb_\mu^3 + g_1(y_1 - y_R)a_\mu\right)\right]E^0 = \overline{\nu}_e(\alpha)\gamma_\mu\left(Q_\nu + Q_\nu^{'}\gamma_5\right)\nu_e(\alpha)A_\mu + \overline{e}^-\gamma_\mu\left(Q_{oe} + Q_{oe}^{'}\gamma_5\right)e^- A_\mu + \\ &+ \overline{E}^0\gamma_\mu\left(Q_{E^0} + Q_{E^0}^{'}\gamma_5\right)E^0 A_\mu + \overline{\nu}_e(\alpha)\gamma_\mu\left(g_V^\nu + g_A^\nu\gamma_5\right)\nu_e(\alpha)Z_\mu + \overline{e}^-\gamma_\mu\left(G_V^e + G_A^e\gamma_5\right)e^- Z_\mu + \overline{E}^0\gamma_\mu\left(G_V^{E^0} + G_A^{E^0}\gamma_5\right)E^0 Z_\mu \end{aligned}$$

were

$$O_\mu = \gamma_\mu(1+\gamma_5), \qquad O'_\mu = \gamma_\mu(1-\gamma_5),$$

$$W^{\pm}_\mu = \frac{1}{\sqrt{2}}(b^1_\mu \mp i b^2_\mu), \quad \hat{\partial} = \gamma_\mu \partial_\mu, \ \overline{\psi} = \psi^+ \gamma_4,$$

$$Q_\nu = \frac{1}{4}\left[g\sin\theta + g_1 y_L \cos\theta\right],$$

$$Q'_{0e} = \frac{1}{4} g_1 (y_L - y_R)\cos\theta,$$

$$Q'_\nu = \frac{1}{4}\left[g\sin\theta + g_1 y_L \cos\theta\right],$$

$$Q_{E^0} = \frac{1}{4}\left[g\sin\theta + g_1(y_1 + y_R)\cos\theta\right], \quad (9a)$$

$$Q_{0e} = \frac{1}{4}\left[-2g\sin\theta + g_1(y_L + y_R)\cos\theta\right],$$

$$Q'_{E^0} = \frac{1}{4}\left[-g\sin\theta + g_1(y_1 - y_R)\cos\theta\right],$$

$$g^\nu_V = \frac{1}{4}\left[g\cos\theta - g_1 y_L \sin\theta\right],$$

$$G^e_A = \frac{1}{4} g_1 (y_R - y_L)\sin\theta,$$

$$g^\nu_A = \frac{1}{4}\left[g\cos\theta - g_1 y_L \sin\theta\right],$$

$$G^{E^0}_V = \frac{1}{4}\left[g\cos\theta - g_1(y_1 + y_R)\sin\theta\right], \quad (9b)$$

$$G^e_V = \frac{1}{4}\left[-2g\cos\theta - g_1(y_L + y_R)\sin\theta\right],$$

$$G^{E^0}_A = \frac{1}{4}\left[-g\cos\theta - g_1(y_L - y_R)\sin\theta\right].$$

From expressions (9a) it is clear that, as in the case of work [12], here too the P-invariance of the electromagnetic interaction and the electroneutrality condition for the neutrino are equivalent.

RESULTS AND DISCUSSION

In the case under consideration, the lepton masses are generated using the Yukawa mass Lagrangian

$$L^l_{mass} = \frac{m_e}{\langle\phi\rangle}\overline{\psi}_R \psi_L \phi + \frac{m_{E^0}\cos\alpha}{\langle\varphi_0\rangle}\overline{\psi}_R \psi'_L \varphi^c + h.c. \quad (10)$$

where $\varphi^c = i\tau_2 \varphi^*$, $\sin\alpha = \dfrac{m_e}{m_{E^0}}$.

Taking into account the conservation of hypercharge, from (10) we obtain

$$y_R = y_L + y_\phi, \ y_R = y_1 - y_\varphi. \quad (11)$$

Taking into account (11) in (9a) we obtain

$$Q'_{E^0} = \frac{1}{4}\left(-g\sin\theta + g_1 y_\varphi \cos\theta\right) \quad (12)$$

If we take into account the P – invariance of the electromagnetic interaction of the new lepton ($Q'_{E^0} = 0$) and the electron ($Q'_{0e} = 0$), from (9a) and (12) we obtain

$$y_L = y_R, \ g_1 y_\varphi \cos\theta = g\sin\theta. \quad (13)$$

Taking into account these relations and (11) in expressions (9a), we obtain

$$Q'_\nu = Q_\nu = \frac{Q_e}{4}\left(1 + \frac{y_L}{y_\varphi}\right), \quad Q_{0e} = -\frac{Q_e}{2}\left(1 - \frac{y_L}{y_\varphi}\right),$$

$$Q_{E^0} = \frac{Q_e}{2}\left(1 + \frac{y_L}{y_\varphi}\right) \quad (14)$$

where $Q_e = g\sin\theta = |e|$ is the absolute value of the electron charge.

From (14), it is clear that the electric charges of the leptons depend on the hypercharge of the Higgs field, and this can be seen as evidence of quantization of the electric charge of the leptons. If we take into account the P-invariance of the electromagnetic interaction of neutrinos ($Q'_\nu = 0$) from (14), (13), and (11), we obtain

$$y_L = y_R = -y_\varphi, \ y_\phi = 0. \quad (15)$$

If in (9b), we take into account relations (11), (13), and (15), then for the vector and axial constants of the interactions of leptons with the Z boson, we obtain

$$g^\nu_V = g^\nu_A = \frac{g}{4\cos\theta},$$

$$G^e_V = \frac{g}{4\cos\theta}\left(-2 + 4\sin^2\theta\right), \qquad G^e_A = 0,$$

$$G^{E^0}_V = -G^{E^0}_A = \frac{g}{4\cos\theta} \quad (16).$$

From expression (16) it is evident that the constants of interactions of leptons with the Z boson coincide with the expressions in the work [15] and when obtaining them for the hypercharge of particles, specific numbers were not taken.

Taking into account the covariant derivatives D^φ_μ , D^ϕ_μ from (4) and moving on to the vacuum expectation values of the fields (2)

$$\langle\varphi\rangle=\begin{pmatrix}0\\ \langle\varphi_0\rangle\end{pmatrix}=\frac{1}{\sqrt{2}}\begin{pmatrix}0\\ \eta\end{pmatrix},\qquad \langle\phi\rangle=\frac{\upsilon}{\sqrt{2}}$$

we obtain the following expression for the part of the Lagrangian responsible for the masses of the vector bosons:

$$\left(D_\mu^\varphi\varphi\right)^+\left(D_\mu^\varphi\varphi\right)+\left(D_\mu^\phi\phi\right)^+\left(D_\mu\phi\right)=M_W^2W_\mu^+W_\mu^-+\frac{1}{2}M_Z^2Z_\mu^2+\frac{1}{2}M_A^2A_\mu^2+M_ZM_AA_\mu Z_\mu \tag{17}$$

where

$$M_W^2=\frac{g^2\langle\varphi_0\rangle^2}{2},\ M_Z^2=\frac{1}{2}\left[\langle\varphi_0\rangle^2\left(g\cos\theta+g_1y_\varphi\sin\theta\right)^2+\langle\phi\rangle^2\left(g_1y_\phi\sin\theta\right)^2\right],$$

$$M_A^2=\frac{1}{2}\left[\langle\varphi_0\rangle^2\left(g\sin\theta-g_1y_\varphi\cos\theta\right)^2+\langle\phi\rangle^2\left(g_1y_\phi\cos\theta\right)^2\right]$$

the mass terms of the fields $W_\mu^\pm$, Z_μ and A_μ.

If we take into account relations (13) and (15) in expressions (17), then for the mass of the vector fields we have

$$M_W=\frac{g\langle\varphi_0\rangle}{\sqrt{2}},\qquad M_A=0,$$

$$M_Z=\frac{M_W}{\cos\theta}\quad .$$

CONCLUSIONS

Thus, from the P-invariance of the electromagnetic interactions of the neutrino $\left(Q_\nu^{'}=0\right)$, electron $\left(Q_{oe}^{'}=0\right)$ and E^0 lepton $\left(Q_{E^0}^{'}=0\right)$ it follows that the photon mass is zero, and between the masses of the $W_\mu^\pm$ and Z - bosons there is a well-known relation $M_W=M_Z\cos\theta$.

Considering relation (15) in expressions (14), for the charge of the leptons we have

$$Q_\nu=0\,,Q_{E^0}=0\,,Q_{0e}=-Q_e=-|e|\,. \tag{18}$$

In conclusion, we note that the dependence of the electric charge of leptons on the hypercharge of the Higgs field (expression (14)) and the fixation of the hypercharge of lepton isodoublets by the hypercharge of the Higgs field (relation $y_L=y_R=-y_\varphi$) can be interpreted as the influence of the Higgs field on the quantization of the electric charge of leptons. This means that it can be assumed that Higgs fields, interacting with particles, give them mass and also play a role in the "formation" of electric charge.

[1] *C.Q. Geng.* Remarks on charge quantization of fermions and bosons, Phys.Rev. D, 41(1990) 1292-1295.

[2] *S. Rudaz.* Electric-charge quantization in the standard model, Phys.Rev. D , 41 (1990) 2619-2621.

[3] *J. Sadkowski, M. Zraek.* Charge quantization in the standard model with three generations of fermions, Phys.Rev.D, 45(1992) 1701-1707.

[4] *O.B. Abdinov, F.T. Khalil-zade, S.S. Rzaeva.* $SU(3)_C$x $SU(3)_L$x U(1)xU(1) model of electroweak interaction and electric charge quantization , AJP Physics , XV (2009) 24–33.

[5] *O.B. Abdinov, F.T. Khalil-zade, S.S. Rzayeva.* Quantization of electric charge in gauge theories , Письма в ЭЧАЯ , 7 (2010) 516–535 .

[6] *He X.-G.* et al.C harge Quantization in Supersymmetric, Technicolor , and Composite Models.Phys.Rev.D. 1989.v.40.p. 3140.

[7] *K.S. Babu, R.N. Mohapatra.* Is There a Connection between Quantization of Electric Charge and a Majorana Neutrino ? Phys.Rev. Lett. 1989 .v.63. p. 938-941.

[8] *R. Foot, H. Lew, R.R. Volkas.* Phenomenology of Quark-Lepton Symmetric Models. Phys.Rev. D.1991.v.44.p.1531.

[9] *P.V. Dong, H.N. Long.* Electric Charge Quantization in $SU(3)_C$ x$SU(3)_L$ xU(1) Models.hep-ph\0507155 v1.2005.

[10] *O.B. Abdinov, F.T. Khalil-zade, S.S. Rzaeva.* Electric Charge Quantization in Standard Model.hep-ph\0807.4359v1.

[11] K.S. Babu, R.N. Mohapatra .Why does electromaqnetism conserve parity? Phys.Rev. D., 42 (1990)3866-3869.

[12] *B.I. Mekhtiev, F.T. Khalil- zade.* Standard model and quantization electric charge of leptons , Transactions of the National Acadmyyemy of Science of Azerbaijan, series of physicmathematical and technical sciences, Physics and Astronomy , XLIV, No. 5 (2024) 14–18.

[13] *B.I. Mehdiyev, F.T.Khalil-zade*. Quantization of the electric charge of quarks in the standard model, Transactions of the National Academy of Science of Azerbaijan, series of physics, mathematical, and technical sciences, Physics and Astronomy, XLV, no. 2 (2025), 14–18.

[14] *B.I. Mehdiyev, F.T.Khalil- zade*. Quantization of electric charge of leptons in the Weinberg-Salam model with a right-hand neutrino component, AJP Physika, XXXI (2025), 49–52.
[15] *N.A. Quliev, I.Q. Jafarov, F.T. Khalil-zade, B.I. Mehtiyev, R.Sh. Yakhyayev*. The unified SU(2)xU(1)x U' (1) –models of electroweak interaction of leptons, Annalen der Physik, 41(1984)93-108.

Received: 22.12.2025